

TECHNICAL REPORT

**Remediation and Treatment Technology
Development and Support for DOE Oak Ridge
Office: Parameterization of Major Transport
Processes of Mercury Species**

Date submitted:

February 18, 2013

Principal Investigators:

Leonel E. Lagos, Ph.D., PMP®

David Roelant, Ph.D.

Florida International University Collaborators:

Georgio Tachiev, PhD, PE, Project Manager

Yong Cai, PhD

Yanbin Li, PhD

Submitted to:

U.S. Department of Energy
Office of Environmental Management
Under Grant # DE-EM0000598



Applied Research Center

FLORIDA INTERNATIONAL UNIVERSITY

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States government. Neither the United States government nor any agency thereof, nor any of their employees, nor any of its contractors, subcontractors, nor their employees makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe upon privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States government or any other agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States government or any agency thereof.

Table of Contents

1. Introduction.....	4
2. Background.....	4
3. Technical Approach.....	5
3.1. Development of the isotope dilution method for analyzing inorganic Hg concentration in water	5
3.2. Relative importance of thiol group versus other groups in L-cysteine in cinnabar dissolution....	6
3.3. Comparison of reduced glutathione (GSH) and oxidized glutathione(GSSG) in promoting cinnabar dissolution	6
3.4. Development of the isotope dilution (ID)-phenylation-purge and trap-ICP-MS technique for analyzing organomercury species	6
4. Results.....	6
4.1. Development of the isotope dilution method for analyzing inorganic Hg concentration in water	6
4.2. Relative importance of thiol group versus other groups in L-cysteine in cinnabar dissolution....	7
4.1. Comparison of reduced glutathione (GSH) and oxidized glutathione(GSSG) in promoting cinnabar dissolution	8
4.2. Development of the isotope dilution (ID)-phenylation-purge and trap-ICP-MS technique for analyzing organomercury species	8
5. References.....	10

List of Figures

<i>Figure 1. Coupling of a Flow Injection System (FIAS 400) to ICP-MS for analyzing mercury isotopes.....</i>	<i>7</i>
<i>Figure 2. Effect of thiol group on the release of dissolved Hg from cinnabar</i>	<i>8</i>
<i>Figure 3. Typical chromatogram of pool water sample spiked with Me200Hg and Et199Hg and analyzed by aqueous phenylation followed by purge and trap-ICP-MS.....</i>	<i>9</i>

1. Introduction

The overall objective of this task is to provide laboratory measurements for critical mercury (Hg) transport, transformation, and exchange processes (i.e., methylation/demethylation, and dissolution) to be used in the numerical model. The laboratory experimental work will provide insight on parameters relevant to the Oak Ridge Reservation (ORR) and which are required in the numerical model, such as dissolution rate of mercury. In addition, experimental work will be conducted to analyze the effect of significant environmental factors (pH, Eh, thiols) on the major transport and transformation processes of Hg.

Under this task we will systematically investigate the stability, bioavailability, and mobility of the aged mercury species in soils and sediments. New techniques will be developed and applied in the analysis of Hg species and the study of Hg transport and transformation processes. Effects of environmental factors, such as oxidation-reduction, pH, and complexation with ligands (e.g., cysteine, glutathione, and large molecular NOM), on mobilizing cinnabar will be particularly investigated. From June 2012 to January 2013, we 1) developed an isotope dilution method for analyzing inorganic Hg concentration in water; 2) evaluated the relative importance of thiol group versus other groups in L-cysteine in cinnabar dissolution; 3) tested the difference of reduced glutathione (GSH) and oxidized glutathione (GSSG) in promoting cinnabar dissolution; 4) are developing an isotope dilution (ID)-phenylation-purge and trap-ICP-MS technique for analyzing organomercury species; 5) are developing a new technique using isotope tracers to simultaneously determine the dissolution of cinnabar and re-adsorption of released Hg^{2+} on the cinnabar surface.

2. Background

Mercury occurs in a variety of species, such as elemental mercury (Hg^0), mercury sulfide (HgS), mercury (hydr)oxides (e.g., HgO and $\text{Hg}(\text{OH})_2$), ionic mercury salts (e.g., HgCl_2), and methylmercury (MeHg), in the environment. At historically mercury contaminated sites (e.g., the DOE Oak Ridge site), the originally discharged mercury would undergo complicated aging processes in the environment and eventually be sequestered in the soil and sediment in relatively stable forms (e.g. mercury bead and HgS). The mercury species sequestered in soils and sediments is generally considered low in solubility and reactivity, which prevents this mercury from being released into the solution and from being transported into the aquatic environment. Despite the limited solubility, mobility, and reactivity, the aged mercury species in soils and sediments may undergo a series of physical and chemical processes under certain environmental conditions, releasing dissolved mercury species into the solution. Cinnabar has often been suggested to act as an important sink for mercury in soils and sediments because of its extremely low solubility product (Mikac et al., 2002b; Mikac et al., 2003b). However, recent studies suggested that cinnabar can also release mercury to environment, which could be promoted by the oxidization of S(II) by Fe(III)-(hydr)oxides or dissolved oxygen (DO) (Burkstaller et al., 1975; Pyzik and Sommer, 1981; Barnett et al., 2001a; Mikac et al., 2002a; Mikac et al., 2003a;

Poulton et al., 2004; Holley et al., 2007a) and the presence of ligands such as sulfide, polysulfides, NOM (e. g. natural organic matter), and chloride in the environment (Burkstaller et al., 1975; Paquette and Helz, 1995; Paquette and Helz, 1997; Ravichandran et al., 1998b; Waples et al., 2005). The dissolution of cinnabar is an important process of Hg cycling in aquatic environments as it can provide a sufficient and continuous source of inorganic mercury and subsequently enhance the production of a more toxic species of Hg, methylmercury (MeHg).

Several possible mechanisms have been previously proposed for the dissolution of cinnabar in the environment. The most common one is that cinnabar dissolution is enhanced via the formation of HgL_x complexes, especially the formation of Hg-thiol complexes (MAHALINGAM RAVICHANDRAN et al., 1998; Ravichandran et al., 1998a; Aiken et al., 2001; Ravichandran, 2004; Waples et al., 2005). The coordination between dissolved Hg^{2+} and ligands will reduce the concentration of free Hg^{2+} and thus promote the dissolution of cinnabar. Another proposed mechanism is that the oxidation of S^{2-} will decrease the concentration of S^{2-} and thus promote the dissolution of cinnabar (Burkstaller et al., 1975; Barnett et al., 2001b). However, little is known about the relative importance of the oxidation of S^{2-} and the formation of Hg complexes in HgS dissolution. In addition to the dissolution of HgS, adsorption of Hg^{2+} could also play an important role in the release of Hg^{2+} from cinnabar. Several studies have found that Hg^{2+} could be quickly absorbed on cinnabar in DI water (Burkstaller et al., 1975; MAHALINGAM RAVICHANDRAN et al., 1998; Barnett et al., 2001b; Holley et al., 2007b), indicating the released Hg^{2+} from cinnabar could be re-adsorbed on the cinnabar surface. However, this process has been omitted in most recent studies on cinnabar dissolution and there is lack of a technique that can simultaneously measure the dissolution and re-adsorption of released Hg^{2+} . Such a technique is necessary for measuring the 'real' dissolution rate of cinnabar and elucidating the pathways through which cinnabar dissolves in natural environment.

3. Technical Approach

3.1. Development of the isotope dilution method for analyzing inorganic Hg concentration in water

Isotope dilution method is based on the addition of a known amount of enriched isotopes to samples. The alteration of the isotope ratio can be used to calculate the concentration of Hg in samples. Prior to analysis of the water sample, a certain amount of ^{199}Hg will be added to samples, serving as an internal standard. Then, signals of ^{199}Hg , and ^{202}Hg isotopes will be analyzed using flow injection (FIAS 400 from Perkin-Elmer company) coupled with ICP/MS (ELAN DRC-e ICP/MS from Perkin-Elmer company). Concentration of Hg in samples can be calculated according to the ratio of ^{199}Hg and ^{202}Hg .

3.2. Relative importance of thiol group versus other groups in L-cysteine in cinnabar dissolution

Effects of L-serine and L-cysteine on cinnabar dissolution were tested and compared. The initial concentrations of L-cysteine and L-serine were 10 μM . pH was adjusted to 8.0 with 10 mM NaOH. Samples (~1.0 mL) were collected after 24 h of dissolution experiments, filtered by 0.22 μm polyethylene membrane and analyzed for total Hg.

3.3. Comparison of reduced glutathione (GSH) and oxidized glutathione (GSSG) in promoting cinnabar dissolution

Effects of GSH and GSSG on cinnabar dissolution were tested and compared. The initial concentrations of GSH and GSSG were 10 μM . pH was adjusted to 8.0 with 10 mM NaOH. Samples (~1.0 mL) were collected after 24 h of dissolution experiments, filtered by 0.22 μm polyethylene membrane and analyzed for total Hg.

3.4. Development of the isotope dilution (ID)-phenylation-purge and trap-ICP-MS technique for analyzing organomercury species

Prior to analysis of the water samples, a certain amount of Me^{200}Hg and Et^{199}Hg will be added to samples, serving as internal standards. Me^{200}Hg was synthesized from ^{200}HgO by methylcobalamin. Et^{199}Hg was synthesized from ^{199}HgO by $(\text{C}_2\text{H}_5)_4\text{Sn}$. Me^{200}Hg , Me^{202}Hg , Et^{199}Hg , and Et^{202}Hg were then detected using phenylation-purge trap-ICP-MS. Concentrations of MeHg and EtHg in samples were then calculated according to the ratios of $\text{Me}^{200}\text{Hg}/\text{Me}^{202}\text{Hg}$ and $\text{Et}^{199}\text{Hg}/\text{Et}^{202}\text{Hg}$.

4. Results

4.1. Development of the isotope dilution method for analyzing inorganic Hg concentration in water

- *Coupling of a Flow Injection System (FIAS 400) to ICP-MS for analyzing mercury isotopes*

As shown in Figure 1, a Flow Injection System (FIAS 400) was successfully coupled to ICP-MS and it could be used to analyze mercury isotopes in samples. Influence of nebulizer gas flow rate (0.5 L/min, 0.6 L/min, 0.8 L/min, 0.96 L/min) on the sensitivity was tested. 0.8L/min was chosen as the flow rate of nebulizer gas.




Figure 1. Coupling of a Flow Injection System (FIAS 400) to ICP-MS for analyzing mercury isotopes

- **Development of the isotope dilution method for analyzing inorganic Hg concentrations in water**

Prior to analysis of the water sample, a certain amount of ^{199}Hg was added to samples to form a final concentration of 10 ppt. Concentrations of Hg^{2+} in samples were then calculated according to Eq. 1. The method detection limit was estimated to be 6.7 ppt.

$$C_s = \frac{C_{199}A_{199}^{202} - RC_{199}A_{199}^{199}}{RA^{199} - A^{202}} \quad (1)$$

R: ratio of $^{202}\text{Hg}^{2+}$ to $^{199}\text{Hg}^{2+}$ in samples

C_s : the concentration of Hg in samples.

C_{199} : concentrations of spiked ^{199}Hg .

: the abundance of ^{199}Hg in ^{199}Hg .

: the abundance of ^{202}Hg in ^{199}Hg .

: the abundance of ^{199}Hg in natural Hg.

: the abundance of ^{202}Hg in natural Hg.

4.2. Relative importance of thiol group versus other groups in L-cysteine in cinnabar dissolution

As shown in Figure 2, Hg^{2+} released from cinnabar in $10\mu\text{M}$ L-cysteine was observed to be 40~60 times higher than that in $10\mu\text{M}$ of L-serine or sodium nitrate control medium (Figure 2), indicating that thiol is the major functional group in L-cysteine that promote the dissolution of cinnabar.




Figure 2. Effect of thiol group on the release of dissolved Hg from cinnabar

4.1. Comparison of reduced glutathione (GSH) and oxidized glutathione (GSSG) in promoting cinnabar dissolution

Concentrations of released Hg (II) from cinnabar were analyzed to be 9.1 ± 1.4 , 11.4 ± 3.1 and $116.6 \pm 1.7 \mu\text{g L}^{-1}$ in the presence of in $10 \mu\text{M}$ of GSSG, GSH and L-cysteine, respectively. Total concentration of Hg released from cinnabar in the presence of GSH was only about 10% of that in the presence of L-cysteine. Hg^{2+} released in the presence of GSSG was slightly less than that in the presence of GSH (Figure 2).

4.2. Development of the isotope dilution (ID)-phenylation-purge and trap-ICP-MS technique for analyzing organomercury species

$100 \mu\text{l}$ $167.8 \text{ ng/L Me}^{200}\text{Hg}$ and $100 \mu\text{l}$ $168.6 \text{ ng/L Me}^{199}\text{Hg}$ was added to water samples. Me^{202}Hg , Me^{200}Hg , Me^{199}Hg , Me^{201}Hg , Et^{202}Hg , Et^{200}Hg , Et^{199}Hg , and Et^{201}Hg were detected using phenylation-purge and trap-ICP-MS technique (Figure 3). Concentrations of MeHg and EtHg in water sample could be calculated by the ratio of $\text{Me}^{202}\text{Hg}/\text{Me}^{200}\text{Hg}$ according to Eq. 2. and $\text{Et}^{202}\text{Hg}/\text{Et}^{199}\text{Hg}$ (Eq. 3), respectively. The concentrations of MeHg and EtHg (spiked) in pool waters were measured to be $0.11 \pm 0.004 \text{ ng/L}$ and $0.20 \pm 0.002 \text{ ng/L}$, with a good reproducibility (RSD < 5%). The developed method was then validated by analyzing MeHg and EtHg (spiked) in reference sediment (IAEA 405) and fish materials (DORM 4). The recoveries of EtHg in IAEA 405 and DORM 4 were $104.1 \pm 3.3\%$ for IAEA 405 and $99.7 \pm 2.7\%$ for DORM 4. The recovery of MeHg was $103.3 \pm 3.2\%$ and $106.9 \pm 0.98\%$, respectively. RSDs of the measured concentrations of MeHg and EtHg in sediment and fish samples were also lower than 5%, indicating the high precision of the developed method.

$$C_s = \frac{C_{200}A_{200}^{202} - RC_{200}A_{200}^{200}}{RA^{200} - A^{202}} \quad (2)$$

R: ratio of Me²⁰²Hg to Me²⁰⁰Hg in samples

C_s: the concentration of MeHg in samples.

C₂₀₀: concentrations of spiked Me²⁰⁰Hg.

: the abundance of ²⁰⁰Hg in Me²⁰⁰Hg.

: the abundance of ²⁰²Hg in Me²⁰⁰Hg.

: the abundance of ²⁰⁰Hg in natural Hg.

: the abundance of ²⁰²Hg in natural Hg.

$$C_s = \frac{C_{199}A_{199}^{202} - RC_{199}A_{199}^{199}}{RA^{199} - A^{202}} \quad (3)$$

R: ratio of Et²⁰²Hg to Et¹⁹⁹Hg in samples

C_s: the concentration of EtHg in samples.

C₁₉₉: concentrations of spiked Et¹⁹⁹Hg.

: the abundance of ¹⁹⁹Hg in Et¹⁹⁹Hg.

: the abundance of ¹⁹⁹Hg in Et¹⁹⁹Hg.

: the abundance of ¹⁹⁹Hg in natural Hg.

: the abundance of ²⁰²Hg in natural Hg.




Figure 3. Typical chromatogram of pool water sample spiked with Me200Hg and Et199Hg and analyzed by aqueous phenylation followed by purge and trap-ICP-MS

5. References

- Aiken, G., Waples, J., Nagy, K.L., Ryan, J., Ravichandran, M., 2001. Dissolution of cinnabar by dissolved organic matter. Abstracts of Papers of the American Chemical Society 222, U425-U425.
- Barnett, M.O., Turner, R.R., Singer, P.C., 2001a. Oxidative dissolution of metacinnabar (b-HgS) by dissolved oxygen. *Appl. Geochem.* 16, 1499.
- Barnett, M.O., Turner, R.R., Singer, P.C., 2001b. Oxidative dissolution of metacinnabar (β -HgS) by dissolved oxygen. *Applied Geochemistry* 16, 1499-1512.
- Burkstaller, J.E., McCarty, P.L., Parks, G.A., 1975. Oxidation of cinnabar by iron(III) in acid mine waters. *Environmental Science & Technology* 9, 676-678.
- Holley, E.A., James McQuillan, A., Craw, D., Kim, J.P., Sander, S.G., 2007a. Mercury mobilization by oxidative dissolution of cinnabar ($[\alpha]$ -HgS) and metacinnabar ($[\beta]$ -HgS). *Chemical Geology* 240, 313-325.
- Holley, E.A., James McQuillan, A., Craw, D., Kim, J.P., Sander, S.G., 2007b. Mercury mobilization by oxidative dissolution of cinnabar (α -HgS) and metacinnabar (β -HgS). *Chemical Geology* 240, 313-325.
- MAHALINGAM RAVICHANDRAN , GEORGER . AIKEN , MICHAELM . REDDY , RYAN, J., 1998. Enhanced Dissolution of Cinnabar (Mercuric Sulfide) by Dissolved Organic Matter Isolated from the Florida Everglades. *Environ. Sci. Technol.*, 6.
- Mikac, N., Foucher, D., Niessen, S., Fischer, J.-C., 2002a. Extractability of HgS (cinnabar and metacinnabar) by hydrochloric acid. *Analytical and Bioanalytical Chemistry* 374, 1028-1033.
- Mikac, N., Foucher, D., Niessen, S., Fischer, J.C., 2002b. Extractability of HgS (cinnabar and metacinnabar) by hydrochloric acid. *Analytical and Bioanalytical Chemistry* 374, 1028-1033.
- Mikac, N., Foucher, D., Niessen, S., Lojen, S., Fischer, J., 2003a. Influence of chloride and sediment matrix on the extractability of HgS (cinnabar and metacinnabar) by nitric acid. *Anal. Bioanal. Chem.* 377, 1196.
- Mikac, N., Foucher, D., Niessen, S., Lojen, S., Fischer, J.C., 2003b. Influence of chloride and sediment matrix on the extractability of HgS (cinnabar and metacinnabar) by nitric acid. *Analytical and Bioanalytical Chemistry* 377, 1196-1201.
- Paquette, K., Helz, G., 1995. SOLUBILITY OF CINNABAR (RED HGS) AND IMPLICATIONS FOR MERCURY SPECIATION IN SULFIDIC WATERS. *Water Air and Soil Pollution* 80, 1053-1056.
- Paquette, K.E., Helz, G.R., 1997. Inorganic speciation of mercury in sulfidic waters: The importance of zero-valent sulfur. *Environmental Science & Technology* 31, 2148-2153.
- Poulton, S.W., Krom, M.D., Raiswell, R., 2004. A revised scheme for the reactivity of iron (oxyhydr)oxide minerals towards dissolved sulfide. *Geochimica et Cosmochimica Acta* 68, 3703-3715.
- Pyzik, A.J., Sommer, S.E., 1981. Sedimentary iron monosulfides: Kinetics and mechanism of formation. *Geochimica et Cosmochimica Acta* 45, 687-698.
- Ravichandran, M., 2004. Interactions between mercury and dissolved organic matter—a review. *Chemosphere* 55, 319-331.
- Ravichandran, M., Aiken, G.R., Reddy, M.M., Ryan, J.N., 1998a. Enhanced dissolution of cinnabar (mercuric sulfide) by aquatic humic substances. Abstracts of Papers of the American Chemical Society 216, U785-U785.

- Ravichandran, M., Aiken, G.R., Reddy, M.M., Ryan, J.N., 1998b. Enhanced dissolution of cinnabar (mercuric sulfide) by dissolved organic matter isolated from the Florida Everglades. *Environmental Science & Technology* 32, 3305-3311.
- Waples, J.S., Nagy, K.L., Aiken, G.R., Ryan, J.N., 2005. Dissolution of cinnabar (HgS) in the presence of natural organic matter. *Geochimica Et Cosmochimica Acta* 69, 1575-1588.